

PES1 promotes breast cancer by differentially regulating ER α and ER β

Long Cheng, ... , Xiaojie Xu, Qinong Ye

J Clin Invest. 2012;122(8):2857-2870. <https://doi.org/10.1172/JCI62676>.

Research Article

The initiation of breast cancer is associated with increased expression of tumor-promoting estrogen receptor α (ER α) protein and decreased expression of tumor-suppressive ER β protein. However, the mechanism underlying this process is unknown. Here we show that PES1 (also known as Pescadillo), an estrogen-inducible protein that is overexpressed in breast cancer, can regulate the balance between ER α and ER β . We found that PES1 modulated many estrogen-responsive genes by enhancing the transcriptional activity of ER α while inhibiting transcriptional activity of ER β . Consistent with this regulation of ER α and ER β transcriptional activity, PES1 increased the stability of the ER α protein and decreased that of ER β through the ubiquitin-proteasome pathway, mediated by the carboxyl terminus of Hsc70-interacting protein (CHIP). Moreover, PES1 transformed normal human mammary epithelial cells and was required for estrogen-induced breast tumor growth in nude mice. Further analysis of clinical samples showed that expression of PES1 correlated positively with ER α expression and negatively with ER β expression and predicted good clinical outcome in breast cancer. Our data demonstrate that PES1 contributes to breast tumor growth through regulating the balance between ER α and ER β and may be a better target for the development of drugs that selectively regulate ER α and ER β activities.

Find the latest version:

<https://jci.me/62676/pdf>





PES1 promotes breast cancer by differentially regulating ER α and ER β

Long Cheng,^{1,2} Jieping Li,^{1,3} Yongjian Han,¹ Jing Lin,⁴ Chang Niu,¹ Zhichao Zhou,¹ Bin Yuan,¹ Ke Huang,⁵ Jiezhi Li,¹ Kai Jiang,¹ Hao Zhang,¹ Lihua Ding,¹ Xiaojie Xu,¹ and Qinong Ye^{1,2}

¹Department of Medical Molecular Biology, Beijing Institute of Biotechnology, Beijing, People's Republic of China. ²Institute of Cancer Stem Cell, Cancer Center, Dalian Medical University, Liaoning, People's Republic of China. ³Department of Clinical Laboratory, Fuzhou General Hospital of Fujian Corps, CAPF, Fuzhou, People's Republic of China. ⁴Department of Clinical Laboratory, First Affiliated Hospital, and ⁵Department of Obstetrics and Gynecology, Chinese PLA General Hospital, Beijing, People's Republic of China.

The initiation of breast cancer is associated with increased expression of tumor-promoting estrogen receptor α (ER α) protein and decreased expression of tumor-suppressive ER β protein. However, the mechanism underlying this process is unknown. Here we show that PES1 (also known as Pescadillo), an estrogen-inducible protein that is overexpressed in breast cancer, can regulate the balance between ER α and ER β . We found that PES1 modulated many estrogen-responsive genes by enhancing the transcriptional activity of ER α while inhibiting transcriptional activity of ER β . Consistent with this regulation of ER α and ER β transcriptional activity, PES1 increased the stability of the ER α protein and decreased that of ER β through the ubiquitin-proteasome pathway, mediated by the carboxyl terminus of Hsc70-interacting protein (CHIP). Moreover, PES1 transformed normal human mammary epithelial cells and was required for estrogen-induced breast tumor growth in nude mice. Further analysis of clinical samples showed that expression of PES1 correlated positively with ER α expression and negatively with ER β expression and predicted good clinical outcome in breast cancer. Our data demonstrate that PES1 contributes to breast tumor growth through regulating the balance between ER α and ER β and may be a better target for the development of drugs that selectively regulate ER α and ER β activities.

Introduction

The association between estrogen and breast cancer was recognized over 100 years ago. Estrogen exerts its function through its 2 nuclear receptors, estrogen receptor α (ER α) and ER β (1, 2). ER belongs to a superfamily of ligand-activated transcription factors that share structural similarity characterized by several functional domains. N-terminal estrogen-independent and C-terminal estrogen-dependent activation function domains (AF1 and AF2, respectively) contribute to the transcriptional activity of the 2 receptors. The DNA-binding domain of the ERs is centrally located. The ligand-binding domain, overlapping AF2, shows 58% homology between ER α and ER β . The DNA-binding domain is identical between the 2 receptors, except for 3 amino acids. However, the AF1 domain of ER β has only 28% homology with that of ER α . The binding of estrogen to ER leads to ER dimerization and its recruitment to the estrogen-responsive elements (EREs) on the promoters of ER target genes, thereby either enhancing or repressing gene activation.

The development of breast cancer is associated with dysregulation of ER expression (3–8). Compared with that in normal breast tissues, the proportion of cells expressing ER α is increased, whereas ER β expression is reduced, in hormone-dependent breast tumors. The ratio of ER α /ER β expression is higher in breast tumors than in normal tissues, and ER α and ER β are antagonistic to each other. ER α mediates the tumor-promoting effects of estrogens, whereas ER β inhibits breast cancer cell growth. ER β reduces cell proliferation induced by ER α activation. Although ER α and ER β have been shown to have a yin-yang relationship in breast tumorigenesis, the molecular mechanism underlying this process remains unclear.

In this study, we show that PES1 (also known as Pescadillo) plays an essential role in estrogen-induced breast tumor growth through regulation of the yin-yang balance between ER α and ER β and is the first such gene to be identified to our knowledge. PES1, a breast cancer-associated gene 1 (BRCA1) C-terminal (BRCT) domain-containing protein, is estrogen inducible, and its expression gradually increases during breast cancer development and progression (9–11). Theoretically, in the treatment of patients with ER α -positive breast cancer, in which ER β is antagonistic to ER α , a drug that decreases transcriptional activity of ER α but increases that of ER β should be better than the currently used endocrine drugs tamoxifen or fulvestrant, which decrease both ER α and ER β transactivation (12, 13). We show that, through the ubiquitin-proteasome pathway, PES1 enhances ER α levels but reduces ER β protein levels, correlating with their respective physiological activities in breast cancer. Thus, PES1 may represent a very promising target for the development of better drugs for breast cancer endocrine therapy.

Results

PES1 differentially regulates transcriptional activity of ER α and ER β as well as their target genes. To define the exact role of PES1 in breast tumor growth, we investigated whether PES1 regulates estrogen signaling. PES1 overexpression in ER α - and ER β -positive MCF7 cells (Figure 1A), ER α -positive and ER β -negative ZR75-1 and T47D cells (Supplemental Figure 1, A and B; supplemental material available online with this article; doi:10.1172/JCI62676DS1), and ER α - and ER β -negative SKBR3 (Figure 1B) breast cancer cells increased transcription of a luciferase reporter construct containing the ERE in response to the ER α -specific agonist propylpyrazole triol (PPT) but decreased ERE reporter transcription in response to the ER β -specific agonist diarylpropionitrile (DPN). This effect was PES1 specific because expression of the known

Authorship note: Long Cheng and Jieping Li contributed equally to this work.

Conflict of interest: The authors have declared that no conflict of interest exists.

Citation for this article: *J Clin Invest.* 2012;122(8):2857–2870. doi:10.1172/JCI62676.




Figure 1

PES1 differentially regulates transcriptional activity of ER α and ER β and expression of their target genes. (A and B) Luciferase reporter assays of ER α and ER β transcriptional activity in (A) MCF7 or (B) SKBR3 cells transiently transfected with ERE-LUC and PES1, SRC1, GRIP1, XRCC1, or BARD1 with or without ER α or ER β and 24-hour treatment with 10 nM E2, 1 nM PPT, or 1 nM DPN. Results shown are mean \pm SD of 3 independent experiments. $\dagger P < 0.01$, $* P < 0.01$, $\# P < 0.01$, $\ddagger P < 0.01$ versus empty vector in the (A) absence or (B) presence of ER α or ER β with vehicle (-), E2, PPT, and DPN, respectively. (C) Luciferase reporter assays in MCF7 cells stably transfected with PES1 siRNA or PES1-R plus siRNA-resistant PES1 (PES1-R) and treated as above. Immunoblot analysis of PES1 expression is shown. Results shown are mean \pm SD of 3 independent experiments. $\dagger P < 0.01$, $\# P < 0.01$, $\ddagger P < 0.01$ versus control siRNA with E2, PPT, and DPN, respectively. (D) Real-time RT-PCR analysis of 47 genes identified by cDNA microarray in our study and 4 genes identified in other studies (*CCND1*, *CTSD*, *E2F1*, and *C-FOS*) in PES1 knockdown MCF7 cells treated or not treated with E2 (+E2 or -E2, respectively) for 24 hours. Data shown are mean \pm SD of triplicate measurements that have been repeated 3 times with similar results. $\dagger P < 0.05$, $\# P < 0.01$ versus control siRNA without E2. $\ddagger P < 0.05$, $\ddagger P < 0.01$ versus control siRNA with E2. (E) Immunoblot analysis of estrogen-responsive gene expression in PES1 knockdown MCF7 cells.



ER cofactors, steroid receptor coactivator-1 (SRC1) or glutamate receptor-interacting protein 1 (GRIP1), did not oppositely regulate ER α and ER β transactivation, and other BRCT domain-containing proteins, X-ray repair complementing defective repair in Chinese hamster cells 1 (XRCC1) and BRCA1-associated RING domain protein 1 (BARD1), had no effect. As expected, SRC1 and GRIP1 increased the transcriptional activity of both ER α and ER β (Figure 1A). In contrast, siRNA knockdown of endogenous PES1 reduced transcriptional activity of ER α and enhanced that of ER β in MCF7 cells (Figure 1C). These effects could be rescued by PES1 reexpression in PES1 knockdown MCF7 cells.

Next, we performed cDNA microarray analysis to monitor gene expression profiles in PES1 knockdown MCF7 cells. In the presence of 17 β -estradiol (E2), PES1 regulated the expression of 256 genes, including over 127 previously reported E2-regulated genes (refs. 14–22, Supplemental Figure 2, and Supplemental Tables 1–3). Real-time RT-PCR confirmed the PES1-mediated expression of 47 genes identified in our study and 4 well-known E2-regulated genes (cyclin D1 [CCND1], cathepsin D [CTSD], E2F transcription factor 1 [E2F1], and C-FOS) identified in other studies (refs. 14–22 and Figure 1D). The expression of many estrogen-responsive genes known to have important functions in DNA replication (23, 24) and cell cycle regulation (24) was found to be downregulated by PES1 knockdown, including replication factor C (RFC), minichromosome maintenance genes, proliferating cell nuclear antigen (PCNA), E2F1, E2F2, MYB, MYC, cyclin-dependent kinase 1 (CDC2), CCND1, and survivin (BIRC5). Interestingly, some of these genes, such as MYC, CCND1, PCNA, E2F1, and BIRC5, were reported to be activated by ER α but to be repressed by ER β (14–22). Consistent with the results of PES1 knockdown, PES1 overexpression increased the transcription of E2F1, CCND1, cyclin E2 (CCNE2), and CTSD in the presence and/or absence of E2, with higher magnitude in the presence of E2 (Supplemental Figure 3A). The expression of 10 representative estrogen-responsive genes was further confirmed by immunoblotting (Figure 1E).

PES1 differentially regulates the dimerization and promoter occupancy of ER α and ER β . Dimerization is a regulatory mechanism of controlling transcription factor activity. Upon dimerization, transcription factors bind to promoter sequences of target genes. ER α and ER β homodimerization is thought to be critical for ER α and ER β transcriptional activity, whereas ER α -ER β heterodimerization facilitates inhibition of ER α transactivation by ER β (25). In agreement with the findings that ER α and ER β transcriptional activity was oppositely regulated by PES1, overexpression of PES1 increased ER α homodimerization (Figure 2A) and decreased ER β homodimerization and ER α -ER β heterodimerization (Figure 2B). Like ER α and ER β , PES1 was recruited to the estrogen-responsive CTSD, CCND1, E2F1, and CCNE2 promoters but not to an unrelated β -actin promoter (Figure 2C). In addition, unlike ER α and ER β (26, 27), PES1 was not recruited to the distal enhancers of CTSD and CCND1 (Figure 2C). Importantly, consistent with the results of ER α and ER β transactivation, which was oppositely regulated by PES1, PES1 knockdown decreased ER α promoter occupancy but increased that of ER β (Figure 2D).

PES1 has been shown to directly interact with the cadmium response element (CdRE) of the heme oxygenase-1 (HO1) promoter (28). We searched potential CdREs of the CTSD, CCND1, E2F1, and CCNE2 genes and found that CCND1 and E2F1 had putative CdREs. The results of EMSA demonstrated that PES1 bound indeed to the CdRE of the HO1 promoter (Figure 2E). However,

PES1 did not bind to the putative CdREs of CCND1 and E2F1, suggesting that PES1 may regulate estrogen-responsive gene transcription through EREs.

PES1 oppositely modulates ER α and ER β protein stability. To investigate how PES1 increases transactivation of ER α but decreases that of ER β , we examined the effects of PES1 on ER α and ER β expression. In the absence or presence of E2, PES1 knockdown reduced ER α protein expression but enhanced ER β protein levels in MCF7 cells (Figure 3A). Reexpression of PES1 in the knockdown cells rescued this effect. PES1 knockdown did not alter ER α and ER β mRNA levels in MCF7 cells (Supplemental Figure 3B). Similar trends were obtained in ZR75-1 cells and normal human mammary epithelial cells (HMECs) (Supplemental Figure 3, C–F), suggesting that PES1 regulates ER α and ER β expression at the posttranscriptional level.

Recent studies show that at least 3 ER β isoforms, including the wild-type ER β (ER β 1), ER β 2, and ER β 5, are expressed in breast cancer (29, 30). ER β is the only fully functional isoform. ER β 2 and ER β 5 can not bind estrogen. All 3 ER β isoforms can inhibit ER α transcriptional activity (31). Western blot analysis with anti-ER β or anti-MYC showed that ER β 1/ER β was indeed expressed in MCF7 cells, because the location of the endogenous band was similar to that of MYC-tagged ER β 1 but not MYC-tagged ER β 2 and ER β 5 (Supplemental Figure 4A). The anti-ER β used recognized all 3 ER β isoforms (Supplemental Figure 4A, right panel). In addition, PES1 downregulated ER β , ER β 2, and ER β 5 (Supplemental Figure 4B).

Since PES1 modulates ER α and ER β expression at the posttranscriptional level, we first determined the half-life of ER α protein in PES1 knockdown MCF7 cells. In the absence or presence of E2, the half-life of ER α protein in PES1 knockdown cells was reduced from more than 12 hours to approximately 4 hours and from 3 to 2 hours, respectively (Figure 3B). Because ER β is usually expressed at low levels in breast cancer cell lines, we determined the half-life of ER β by ER β overexpression in HEK293T cells. In the absence and presence of E2, PES1 overexpression decreased the half-life of ER β from approximately 12 to 6 hours and from 9 to 3 hours, respectively (Figure 3C).

Next, we determined effects of PES1 on the protein levels of ER α and ER β domains. In transfected HEK293T cells, PES1 increased the protein levels of the ER α AF2 domain but reduced the levels of the ER β AF2 domain (Supplemental Figure 5, A and B). Domain-swapping experiments in which the ER α AF2 domain was exchanged with the ER β AF2 domain abolished the ability of PES1 to regulate ER α and ER β protein levels (Supplemental Figure 5C), indicating important roles of the AF2 domains of ER α and ER β in the regulation of ER α and ER β protein levels by PES1.

To define the region of PES1 that modulates ER α and ER β protein levels, we transfected HEK293T cells with a series of PES1 deletion mutants. The 221–322 region of PES1 (PES1_{A221–322}), but not other regions tested, enhanced ER α protein expression, whereas the 311–588 region of PES1 (PES1_{A311–415}) decreased ER β protein levels (Supplemental Figure 5, D and E). As expected, PES1_{A221–322} and PES1_{A311–415} did not change the expression of ER α and ER β , respectively, but PES1_{A221–322} reduced ER β expression, and PES1_{A311–415} increased ER α expression (Figure 3, D and E). In MCF7 cells, PES1_{A221–322} decreased ER β transcriptional activity, and PES1_{A311–415} increased ER α transcriptional activity, suggesting that the alteration of ER α and ER β protein levels by PES1_{A221–322} and PES1_{A311–415} correlates with their effects on ER α and ER β transactivation (Figure 3F).




Figure 2

PES1 differentially modulates the dimerization and promoter occupancy of ER α and ER β . **(A and B)** Coimmunoprecipitation analysis of ER α and ER β homodimerization and ER α -ER β heterodimerization. HEK293T cells transiently transfected with the indicated HA-, FLAG-, and MYC-tagged constructs were immunoprecipitated with **(A)** anti-FLAG or **(B)** anti-MYC antibodies, followed by immunoblotting as indicated. **(C)** ChIP analysis of the occupancy of PES1, ER α , or ER β on the indicated estrogen-responsive proximal promoters (PPs) or distal enhancers (DEs) in MCF7 cells treated with 10 nM E2 for 1 hour. The β -actin promoter was included as a negative control. IgG, normal serum. Data shown are mean \pm SD of triplicate measurements that have been repeated 3 times with similar results. * $P < 0.05$, † $P < 0.01$ versus respective IgG without E2. # $P < 0.01$ versus respective IgG with E2. **(D)** ChIP analysis of the occupancy of ER α or ER β on CTSD and CCND1 promoters in MCF7 cells stably transfected with control siRNA or PES1 siRNA and treated as in **C**. Data shown are mean \pm SD of triplicate measurements that have been repeated 3 times with similar results. * $P < 0.01$ versus respective control siRNA without E2. # $P < 0.01$ versus respective control siRNA with E2. **(E)** EMSA using the in vitro-translated PES1 and the biotin-labeled CdRE or mutated CdRE (mCdRE) probe. Cold probe was used for competition experiments.

The E3 ubiquitin ligase CHIP is important for PES1 modulation of ER α and ER β protein stability. The finding that PES1 regulates ER α and ER β protein stability suggests that the ubiquitin-proteasome pathway may be involved in this process. Indeed, addition of the proteasome inhibitor MG132 or lactacystin blocked PES1 knock-down-mediated ER α degradation and PES1 overexpression-mediated

ER β degradation (Figure 4, A and B, and data not shown). Overexpression of PES1 or PES1 $_{\Delta 311-415}$, which increases ER α protein levels, reduced ER α ubiquitination, whereas expression of PES1 $_{\Delta 211-322}$, which does not increase ER α protein levels, did not (Supplemental Figure 6A). Likewise, overexpression of PES1 or PES1 $_{\Delta 211-322}$, which decreases ER β protein levels, increased ER β




Figure 3

Modulation of ER α and ER β stability by PES1 correlates with their respective transcriptional activity. **(A)** Immunoblot analysis of MCF7 cells stably transfected with PES1 siRNA or PES1 siRNA plus siRNA-resistant PES1 and treated with E2 for 24 hours. **(B)** Immunoblot analysis of ER α in MCF7 cells stably transfected with control siRNA or PES1 siRNA at the indicated times after exposure to the protein synthesis inhibitor cycloheximide (20 mg/ml) in the absence or presence of 10 nM E2. Graphs show quantification of immunoblot data. **(C)** Immunoblot analysis of ER β in HEK293T cells transiently transfected with FLAG-tagged ER β (FLAG-ER β) and MYC-tagged PES1 (MYC-PES1) and treated as in **B**. **(B and C)** Data shown are mean \pm SD of 3 independent experiments. **(D and E)** Immunoblot showing **(D)** ER α and **(E)** ER β protein levels in HEK293T cells transiently transfected with ER α or ER β and FLAG-PES1, FLAG-PES1 $\Delta_{221-322}$, or FLAG-PES1 $\Delta_{311-415}$. **(F)** Luciferase reporter assays of ER α and ER β transcriptional activity in MCF-7 cells transiently transfected with ERE-LUC and FLAG-tagged PES1, PES1 $\Delta_{221-322}$, or PES1 $\Delta_{311-415}$ and treated with 10 nM E2, 1 nM PPT, or 1 nM DPN for 24 hours. Results shown are mean \pm SD of 3 independent experiments. * P < 0.01, # P < 0.01, † P < 0.01 versus empty vector with E2, PPT, and DPN, respectively. Immunoblot analysis of FLAG-tagged PES1, PES1 $\Delta_{221-322}$, or PES1 $\Delta_{311-415}$ in the presence of 10 nM E2 is shown.




Figure 4

PES1 oppositely regulates ERα and ERβ stability through the CHIP-mediated ubiquitin-proteasome pathway. (A) Immunoblot analysis of MCF7 cells stably transfected with PES1 siRNA or control siRNA and treated with 10 nM E2 or 10 nM E2 plus the proteasome inhibitor MG132 (10 μM). (B) Immunoblot analysis of HEK293T cells transiently transfected with FLAG-PES1 and MYC-ERβ and treated as in A. (C and D) Ubiquitination of endogenous ERα and ERβ. Cell lysates from MCF-7 cells stably transfected with PES1 siRNA were immunoprecipitated with antibodies specific for (C) ERα or (D) ERβ, followed by immunoblotting with the indicated antibodies. Ub, ubiquitin. (E and F) Effects of PES1 on CHIP-mediated ERα and ERβ ubiquitination. Coimmunoprecipitation was performed in HEK293T cells transiently transfected with the indicated constructs. (G and H) Effects of PES1 on CHIP-mediated degradation of ERα and ERβ. Western blot analysis of MCF-7 cells transfected with (G) CHIP siRNA and PES1 siRNA or with (H) CHIP siRNA and FLAG-PES1 and treated with 10 nM E2 for 24 hours.

ubiquitination, whereas PES1_{Δ311-415}, which does not decrease ERβ protein levels, did not (Supplemental Figure 6B). Importantly, PES1 knockdown increased the ubiquitination of endogenous ERα but decreased the ubiquitination of endogenous ERβ in MCF7 cells (Figure 4, C and D).

ERα and ERβ are substrates of carboxyl terminus of Hsc70-interacting protein (CHIP) (32–34), an E3 ubiquitin ligase. Intriguingly, PES1 or PES1_{Δ311-415}, but not PES1_{Δ221-322}, reduced CHIP-mediated

ERα ubiquitination (Figure 4E). Likewise, PES1 or PES1_{Δ221-322}, but not PES1_{Δ311-415}, increased CHIP-mediated ERβ ubiquitination (Figure 4F). CHIP knockdown greatly inhibited the ability of PES1 to regulate ERα and ERβ ubiquitination (Supplemental Figure 6, C and D). Furthermore, consistent with the ubiquitination results, CHIP knockdown almost abolished the effects of PES1 on ERα and ERβ degradation (Figure 4, G and H). These data suggest that CHIP plays a key role in the modulation of ERα and ERβ protein levels by PES1.




Figure 5





PES1 and CHIP formed a complex with ERβ but not with ERα. (A–C) Coimmunoprecipitation analysis of the endogenous interaction of PES1 with (A) ERα, (B) ERβ, or (C) CHIP. Cell lysates from MCF-7 cells were immunoprecipitated with antibodies specific for (A) ERα, (B) ERβ, or (C) CHIP, followed by immunoblotting with the indicated antibodies. (D) PES1, ERβ, and CHIP formed a complex. HEK293T cells transfected with the indicated plasmids were immunoprecipitated with anti-FLAG (1st IP). The immune complexes were eluted with FLAG peptide and reimmunoprecipitated (Re-IP) with anti-HA or normal IgG, followed by immunoblotting with the indicated antibodies. (E and F) Effects of PES1 on the interaction of CHIP with (E) ERα and (F) ERβ. Coimmunoprecipitations were carried out in HEK293T cells transiently transfected with the indicated constructs.

PES1 forms a complex with CHIP and ERβ but not with CHIP and ERα. Based on our findings that PES1 regulates CHIP-dependent ERα and ERβ degradation, we tested whether PES1 physically interacts with ER and CHIP. Indeed, GST pull-down and coimmunoprecipitation experiments showed that exogenous PES1 protein associated with exogenous ERα and ERβ proteins as well as exogenous CHIP protein, both in vitro and in vivo (Supplemental Figure 7, A and B, and Supplemental Figure 8, A and B). Importantly, in the absence or presence of E2, endogenous ERα, ERβ, or CHIP from MCF7 cell lysates specifically coimmunoprecipitated with endogenous PES1 (Figure 5, A–C). Moreover, confocal immunofluorescence analysis of MCF7 cells revealed that PES1 also colocalized with ERα and ERβ in the absence or presence of E2 (Supplemental Figure 7, C and D). These data strongly indicate that PES1 interacts with ERα, ERβ, and CHIP.

To map interaction regions of ERα and ERβ in PES1, we performed coimmunoprecipitation experiments using transfected HEK293T cells. PES1₁₋₁₁₀, PES1₁₁₁₋₂₂₀, PES1₂₂₁₋₃₂₂, and PES1₃₁₁₋₅₈₈ interacted with ERα and ERβ, whereas PES1₄₁₅₋₅₈₈ did not (Sup-

plemental Figure 7, E and F). On the other hand, coimmunoprecipitation assays showed that PES1 interacted with the AF1 and AF2 domains, but not the DNA-binding domain, of ERα but only with the AF2 domain of ERβ (Supplemental Figure 7, G and H).

To define the region of CHIP that interacts with ERα, ERβ, and PES1, we used CHIP deletion mutants in coimmunoprecipitation assays. ERβ and PES1 interacted with both CHIP₁₋₁₂₆, containing the tetratricopeptide repeat domain, and CHIP₁₂₇₋₃₀₄, containing the U-box domain, but ERα interacted only with CHIP₁₋₁₂₆ (Supplemental Figure 8, C–E). Importantly, PES1 and CHIP formed a complex with ERβ but not with ERα, possibly because ERβ has more interaction regions in CHIP than ERα (Figure 5D). Based on these observations, we tested whether PES1 affects the interaction of CHIP with ERα and ERβ. Consistent with the effects of PES1 on CHIP-mediated ERα and ERβ degradation, PES1 or PES1_{Δ311-415} reduced the interaction between CHIP and ERα, whereas PES1_{Δ211-322} did not (Figure 5E and Supplemental Figure 9A). Likewise, PES1 or PES1_{Δ211-322} increased the interaction between CHIP and ERβ,



**Figure 6**

PES1 transforms normal HMECs and is required for estrogen-induced breast carcinogenesis. **(A)** Anchorage-dependent growth assays in MCF-7 cells transiently transfected with FLAG-tagged PES1, PES1_{A221-322}, or PES1_{A311-415}. The transfection efficiency is approximately 30%. Cell viability was assessed at the indicated times. * $P < 0.01$ versus empty vector on day 4. # $P < 0.05$, † $P < 0.01$ versus PES1 on day 4. Immunoblot analysis with anti-FLAG is shown. **(B)** Anchorage-dependent growth assays in MCF-7 cells stably transfected with PES1 siRNA or PES1 siRNA plus siRNA-resistant PES1. Cells were treated with 10 nM E2 and analyzed as in **A**. * $P < 0.01$ versus control siRNA without E2 on day 4. # $P < 0.01$ versus control siRNA with E2 on day 4. Immunoblot analysis with anti-PES1 is shown. **(C)** Anchorage-independent growth assays in MCF-7 cells stably transfected as in **B**. Scale bar: 50 μm . **(A–C)** Data are shown as mean \pm SD of 3 independent experiments. * $P < 0.01$ versus control siRNA without E2. # $P < 0.01$ versus control siRNA with E2. **(D)** Volume of xenograft tumors derived from MCF-7 cells expressing control siRNA or PES1 siRNA. Data are shown as mean \pm SD ($n = 8$ for control siRNA; $n = 1$ for PES1 siRNA due to absence of visible tumors in the other 7 mice). * $P < 0.01$ versus control siRNA. Representative tumor tissues were subjected to immunoblot analysis with indicated antibodies. **(E)** Anchorage-independent growth assays in HMECs infected with recombinant lentivirus carrying GFP or PES1. Scale bar: 50 μm . Immunoblotting with the indicated antibodies is shown.

whereas PES1_{A311-415} did not (Figure 5F and Supplemental Figure 9B). These effects are specific for CHIP, because PES1 did not change the interaction of ER α and ER β with E6-associated protein, another E3 ubiquitin ligase for ER α and ER β (refs. 35, 36, and Supplemental Figure 9, C and D).

PES1 is required for estrogen-mediated breast tumor growth. Next, we determined the effect of PES1 on breast cancer cell growth. In assays of anchorage-dependent growth, MCF7 cells transfected with PES1 grew faster than those transfected with PES1_{A211-322}, PES1_{A311-415}, or empty vector, and MCF7 cells transfected with PES1_{A211-322} or PES1_{A311-415} grew faster than those transfected with empty vector (Figure 6A). In contrast, PES1 knockdown almost completely abolished E2-mediated growth stimulation of MCF7 cells (Figure 6B), and this phenotype was rescued by PES1 reexpression. Similar results were observed in ZR75-1 and T47D cells (Supplemental Figure 10, A and B). PES1 knockdown also greatly inhibited anchorage-independent growth of MCF7, ZR75-1, and T47D cells (Figure 6C and Supplemental Figure 10, C and D), and again, the observed effects were rescued by PES1 reexpression in MCF7 cells (Figure 6C). Furthermore, all mice inoculated with MCF7 or ZR75-1 cells expressing control siRNA developed tumors in the presence of E2 but not in the absence of E2 (Figure 6D, Supplemental Figure 10E, and data not shown), suggesting that both MCF7 and ZR75-1 cell lines are estrogen dependent. In contrast, in mice inoculated with MCF7 or ZR75-1 cells expressing PES1 siRNA, only 1 or 3, respectively, out of 8 mice developed tumors in the presence of E2, and these showed late latency and a much smaller tumor size (Figure 6D and Supplemental Figure 10E). The tumors in mice inoculated with MCF7 cells expressing PES1 siRNA had reduced protein levels of ER α , progesterone receptor (PR), MYC, CCND1, and survivin, and increased levels of ER β (Figure 6D). With the exception of those concerning ER β , similar effects were observed in ER β -negative ZR75-1 cells (Supplemental Figure 10E).

Using growth in soft agar as an index of transformation, we examined the effect of PES1 on transformation of HMECs. HMECs with elevated PES1 expression levels, equivalent to those of MCF7 cells, could grow in soft agar, whereas HMECs expressing green fluorescent protein or empty vector could not (Figure 6E). PES1 overexpression increased expression of ER α but decreased that of ER β .

Correlation of PES1 with ER α and ER β in patients with breast cancer. The implication of ER α in breast cancer has been widely investigated with high-quality ER α antibodies. Thus, we examined the specificity of anti-ER β and anti-PES1 antibodies. We confirmed the specificity of the antibodies by immunoblotting of lysates from MCF7 cells transfected with ER β siRNA or PES1 siRNA (Supplemental Figure 11A), immunofluorescence analysis of ER β or PES1 knockdown MCF7 cells (Supplemental Figure 11B), and immunohistochemical staining of breast cancer samples incubated with anti-ER β or anti-PES1 antibodies preincubated with their respective antigens (Supplemental Figure 11C). Intriguingly, consistent with our findings that PES1 upregulated ER α and downregulated ER β s (ER β , ER β 2, and ER β 5), immunohistochemical staining of 116 breast cancer tissues and 92 normal tissues adjacent to breast cancer, using antibodies with confirmed specificity, showed that PES1 expression correlated positively with ER α expression ($P < 10^{-4}$) but negatively with expression of ER β s ($P < 10^{-4}$) (Figure 7, A and B). These data strongly suggest important pathological roles of PES1 in breast cancer.

PES1 predicts clinical outcome of tamoxifen therapy. To determine the clinical relevance of PES1 modulation of ER, we analyzed the survival follow-up information of 65 subjects. PES1 overexpression was significantly associated with better disease-free survival ($P = 0.001$) and overall survival ($P = 0.002$) in patients with breast cancer who received tamoxifen treatment (Figure 7, C and D). To verify the effects of PES1 on tamoxifen sensitivity, we performed animal experiments. Overexpression of PES1 in ZR75-1 cells increased ER α expression and caused increased sensitivity to tamoxifen (Supplemental Figure 10F).

Discussion

An increasing number of studies show that estrogen signaling depends principally on the balance between ER α and ER β expression (3, 4). The disturbance of such balance and, especially, a predominance of proliferative ER α protein over antiproliferative ER β protein may cause cancer in estrogen-responsive organs. Thus, elucidating the regulation of the balance between ER α and ER β expression may not only provide novel mechanistic insights into estrogen-induced tumorigenesis but also improve current endocrine therapy for estrogen-related cancers. Our work reveals for what we believe to be the first time that PES1 plays an essential role in estrogen-induced breast tumor growth through regulation of the yin-yang balance between ER α and ER β (Figure 7E). First, we demonstrated that PES1 enhances transcriptional activity of ER α and reduces that of ER β and modulates many estrogen-responsive genes. Second, consistent with this regulation of ER α and ER β transcriptional activity, PES1 increased the stability of the ER α protein and decreased that of ER β through CHIP-mediated ubiquitin-proteasome pathway. Third, PES1 can transform normal HMECs through increased ER α protein and decreased ER β protein and is required for estrogen-induced breast tumor growth in nude mice. Fourth, expression of PES1 correlates positively with ER α expression and negatively with ER β expression in patients with breast cancer. Since






Figure 7

Association of PES1 with ER α and ER β in breast cancer. (A and B) Expression of ER α , ER β , and PES1 in (A) human breast cancer tissues and (B) normal tissues adjacent to breast cancer. Representative immunohistochemical staining of PES1, ER α , and ER β is shown at top. Original magnification, $\times 20$. Scale bar: 100 μm . A summary of (A) 116 breast cancer tissues or (B) 92 normal breast tissues is shown below, with tissues categorized by low and high expression of PES1 and ER α or ER β . Case 1 and case 2 refer to 2 representative samples. The P value was generated using the χ^2 test. (C and D) Kaplan-Meier estimate of (C) disease-free survival and (D) overall survival in 65 patients with breast cancer who received tamoxifen treatment. Marks on graph lines represent censored samples. High PES1 and low PES1 refer to samples with high and low levels of PES1 expression, respectively. (E) Proposed model for PES1 modulation of the balance between ER α and ER β . The estrogen-inducible protein PES1 blocks interaction of ER α with CHIP through its interaction with CHIP and instead forms a complex with ER β and CHIP, leading to reduced degradation of ER α and increased degradation of ER β by CHIP. In turn, PES1 enhances transcriptional activity of ER α but reduces that of ER β through increased ER α homodimerization and decreased ER β homodimerization and ER α -ER β heterodimerization. Disruption of the balance between ER α and ER β by PES1 contributes to breast tumorigenesis.

dysregulation of the balance between ER α and ER β has also been reported to be associated with other cancers, such as colon cancer and thyroid cancer (37, 38), our data suggest that PES1 may not only act as a determinant of breast tumorigenesis but also play an important role in the development of other cancers. PES1 may be a useful target for hormone-related cancer therapy.

Estrogen stimulates breast cancer cell growth through ER α , and use of antiestrogens blocks this stimulating response (3, 39, 40). As approximately 70% to 80% of all breast cancers are ER α positive at the time of diagnosis, ER α expression has considerable implications for cancer biology and therapy. Tamoxifen is an antiestrogen drug that was developed over 30 years ago and has been widely used in the treatment of all stages of ER α -positive breast cancer (39, 40). Tamoxifen binds and blocks ER α on the surface of cells, preventing estrogens from binding and activating the cell. Due to the discovery of a second form of the ER, ER β , in 1996, the role of tamoxifen in breast cancer endocrine therapy appears to be complicated. Tamoxifen also binds ER β and inhibits ER β transcriptional activity. However, many lines of evidence demonstrate that ER β has an antiproliferative function when reintroduced into ER α -positive breast cancer cells (3, 5–7). In many ways, ER β seems to oppose the action of ER α . The same problem remains for another endocrine drug, fulvestrant (41, 42). It is indicated for the treatment of ER α -positive metastatic breast cancer in postmenopausal women after treatment with other antiestrogens. Fulvestrant binds to ER and prevents structural changes that are necessary for ER to initiate gene transcription. On the other hand, fulvestrant degrades ER, which also reduces gene transcription. Like tamoxifen, fulvestrant also inhibits both ER α and ER β transcriptional activity. Ideally, in the treatment of patients with ER α -positive breast cancer, in which ER β is antagonistic to ER α , a drug that reduces transcriptional activity of ER α but enhances that of ER β should be better than the currently used endocrine drugs tamoxifen or fulvestrant. We demonstrate that PES1 increases ER α but decreases ER β protein levels, correlating with their respective physiological activities in breast cancer. Thus, PES1 represents a very promising target for the development of better drugs for endocrine cancer therapy.

Estrogen has been shown to stimulate the expression of PES1 protein (9–11). Our cDNA microarray analysis indicated that PES1 regulates the expression of 256 genes in MCF7 breast cancer cells, including over 127 previously reported estrogen-responsive genes (14–22). Moreover, PES1 knockdown almost abolishes estrogen-mediated anchorage-dependent and -independent growth of breast cancer cells, and most of mice inoculated with PES1 knockdown breast cancer cells do not develop tumors, even in the presence of estrogen. These results suggest that PES1 is a key mediator of estrogen signaling and that a positive feedback loop of estrogen/PES1 promotes malignant growth of breast cancer cells.

PES1 has been demonstrated to play important roles in embryonic development (43), ribosome biogenesis (44–46), cell cycle regulation (47), and chromosome stability (45), although molecular mechanisms underlying these processes remain largely unknown. We show that PES1 modulates many estrogen-responsive genes that are known to have important functions in DNA replication, cell cycle regulation, and DNA repair. For example, PES1 can regulate the expression of a large number of cell cycle-related genes, including cyclin A2 (*CCNA2*), cyclin B2 (*CCNB2*), *CCND1*, *CCNE2*, cyclin-dependent kinase 1 (*CDK1*), *CDK2*, *E2F1*, *E2F2*, cell-division cycle gene 20 (*CDC20*), *PCNA*, *MYC*, *MYB*, and minichromosome maintenance genes (*MCM2–MCM8* and *MCM10*). In normal human cells, cellular division is an ordered, tightly regulated process, involving multiple cell cycle checkpoints that ensure genomic integrity. Altered regulation of the cell cycle is a hallmark of human cancers (48). Cyclins and their associated CDKs are the central machinery that governs cell cycle progression. Overexpression of cyclin D1, the major regulatory subunit for CDK4, is common in human cancers of epithelial cell origin (49). Approximately 50% of human breast cancers express abnormally high levels of cyclin D1, which is maintained throughout subsequent stages of breast cancer progression, from in situ carcinoma to invasive carcinoma. Both cyclin D1 and CDK2 are required for mammary tumorigenesis induced by the *ErbB-2* oncogene (50). Like cyclin D1, *MYC* and *MYB* are also overexpressed in breast tumors (51, 52). Overexpression of *MYC* contributes to breast cancer development and progression and is associated with poor clinical outcome. *MYC* regulates cell cycle at the G₁/S transition through activation of downstream targets such as cyclin E/CDK2. In addition, *MYC* promotes cell cycle progression through activation of cyclin D1, CDK4, *E2F1*, and *E2F2*. The expression of *MYB* protein was shown to be important for estrogen-stimulated proliferation of breast cancer cells (52). *MYB* controls G₂/M cell cycle transition by direct regulation of cyclin B1 expression. Like cyclin D1, *MYC*, and *MYB*, *E2F1* is also an estrogen-inducible protein (53). *E2F1* is necessary for estrogen regulation of breast cancer cell proliferation. Interestingly, a large number of genes involved in the control of the cell cycle contain regulatory binding sites for *E2F1* (54) (e.g., *CCNA2*, cyclin D3, *CCNE2*, *CDK1*, *MYC*, and *PCNA* as well as *E2F1* itself). DNA replication takes place in the S phase of the cell cycle. The highly orchestrated process of DNA replication ensures the accurate inheritance of genetic information from one cell generation to the next. The MCM proteins are essential for the process of DNA replication (55, 56). Loss of MCM function results in DNA damage and genome instability. The fact that PES1 can regulate many key molecules of cell cycle, DNA replication, and DNA repair suggests the importance of PES1 in breast tumorigenesis and as a therapeutic target. It will be interesting to investigate how PES1 regulates these molecules.



Methods

Plasmids and siRNAs. The estrogen-responsive reporter construct ERE-Luc and eukaryotic expression vectors for FLAG-tagged ER α and ER β have been described previously (57–59). Other mammalian expression vectors encoding FLAG-, MYC-, or HA-fusion proteins tagged at the amino terminus were constructed by inserting PCR-amplified fragments into pcDNA3 (Invitrogen) or pIRESpuro2 (Clontech). Plasmids encoding GST fusion proteins were generated by cloning PCR-amplified sequences into pGEX-KG (Amersham Pharmacia Biotech). The cDNA target sequences of siRNAs for PES1 and CHIP were ACACAAGAAGAAGGTTAAC and GCACGCAAGTACATGGCGGA, respectively, and were inserted into pSilencer2.1-U6neo (Ambion) and pSIH-H1-puro (System Biosciences). Expression vectors for siRNA-resistant PES1 containing a silent mutation in the 3' nucleotide of a codon in the middle of the siRNA-binding site were generated by recombinant PCR. Recombinant lentivirus vectors for PES1 or GFP were made by inserting PCR-amplified fragments into pCDH-EF1-MCS-T2A-puro (System Biosciences).

Cell culture, transfection, and luciferase reporter assay. HEK293T embryonic kidney cells and MCF7, ZR75-1, T47D, and SKBR3 breast cancer cells were routinely cultured in DMEM (Invitrogen) containing 10% FBS (Hyclone). Normal HMECs (Invitrogen) were cultured in HMEC medium (Invitrogen). For hormone treatment experiments, cells were cultured in medium containing phenol red-free DMEM supplemented with 10% charcoal/dextran-treated FBS (Hyclone). Lipofectamine 2000 reagent was used for transfections following the manufacturer's protocol (Invitrogen). Lentiviruses were produced by cotransfection of HEK293T cells with recombinant lentivirus vectors and pPACK Packaging Plasmid Mix (System Biosciences) using Megatran reagent (Origene). Lentiviruses were collected 48 hours after transfection and added to the medium of target cells with 8 μ g/ml polybrene (Sigma-Aldrich). Stable cell lines were selected in 500 μ g/ml G418 or 1 μ g/ml puromycin for approximately 2 months. Pooled clones or individual clones were screened by standard immunoblot protocols and produced similar results. PES1 knockdown stable cell lines grew very slowly and could only be passaged several times. Luciferase reporter assays were performed as described previously (59).

cDNA microarray analysis. cDNA generated from RNA was labeled with Cy3 (PES1 siRNA) and Cy5 (control siRNA), mixed, and hybridized to human oligo chip 35 k v2.0 containing 35,000 human gene elements (CapitalBio). The chip was scanned by LuxScan 10K/A (CapitalBio), and data were analyzed using MAS 3.0 Software (CapitalBio). All results were given as the gene expression ratio (ratio of the intensity of Cy3 to that of Cy5). Genes with more than or equal to 2-fold intensity change were considered of interest and subjected to further investigation by gene ontology analysis and pathway analysis based on the Kyoto Encyclopedia of Genes and Genomes database.

Real-time RT-PCR. Total RNA was isolated using TRIzol Reagent (Invitrogen) and reverse transcribed using SuperScript II Reverse Transcriptase (Invitrogen). Real-time PCR was performed with the primers listed in Supplemental Table 4A as described previously (57).

ChIP assay. ChIP assays were performed as described previously (57) using anti-ER α (Millipore), anti-ER β (Novus Biologicals), and anti-PES1 (Bethyl Laboratories). The primers used for ChIP are listed in Supplemental Table 4B.

Cycloheximide chase assay. Transfected cells were cultured in phenol red-free DMEM medium supplemented with 10% charcoal-stripped FBS for 3 days. Cells were treated with 20 μ g/ml cycloheximide for different time periods in the presence or absence of 10 nM E2. Cell lysates were analyzed by immunoblotting with antibodies specific for ER α (Sigma-Aldrich), FLAG (Sigma-Aldrich), and GAPDH (Santa Cruz Biotechnology Inc.).

Ubiquitination assay. Transfected cells were treated with the proteasome inhibitor MG132 (10 μ M) for 4 hours. The cells were lysed in RIPA buffer and

immunoprecipitated with anti-FLAG (Sigma-Aldrich), anti-ER α (Millipore), or anti-ER β (Novus Biologicals) antibodies as described previously (60). The immunocomplexes were subjected to immunoblot analysis with antibodies specific for MYC or ubiquitin (both from Santa Cruz Biotechnology Inc.).

GST pull-down assay. GST fusion proteins were expressed in pGEX-KG and purified according to the manufacturer's instructions (Amersham Pharmacia Biotech). HEK293T extracts expressing ER α , ER β , or CHIP were mixed with 10 μ g of GST derivatives bound to glutathione-Sepharose beads, and the adsorbed proteins were analyzed as previously described (61).

Coimmunoprecipitation. Cell extracts were prepared, immunoprecipitated, and analyzed as previously described (61). Immunoprecipitation was performed with anti-FLAG M2 Affinity Gel (Sigma-Aldrich), anti-MYC Affinity Gel (Sigma-Aldrich), anti-ER α (Millipore), anti-ER β (Novus Biologicals), or anti-CHIP (Santa Cruz Biotechnology Inc.).

Immunofluorescence. Immunofluorescence was performed as previously described (57). Briefly, cells grown on glass coverslips were fixed, permeabilized, and blocked in normal goat serum. The coverslips were then incubated with rabbit anti-PES1 (Bethyl Laboratories) and mouse anti-ER α (Santa Cruz Biotechnology Inc.) or mouse anti-PES1 (Santa Cruz Biotechnology Inc.) and rabbit anti-ER β (Millipore), followed by incubation with corresponding secondary antibodies. Nuclei were counterstained with DAPI. Confocal images were collected using a Radiance2100 confocal microscope (Bio-Rad).

EMSA. The probes for EMSA were labeled with the Biotin 3'-End DNA Labeling Kit (Pierce) as instructed by the manufacturer. The sequences of the labeled oligonucleotides for the CdRE of the *HO-1* gene (28), the putative CdRE of the *CCND1* gene, and the putative CdRE of the *E2F1* gene were 5'-AATTCGGCGGATTTTGCTAGATTTTGCG-3' (CdRE-*HO-1*), 5'-AGTTTCATATTTGCTAGATATCAGTGTTTG-3' (CdRE-*CCND1*), and 5'-TTTGAACCTGATGCTAGATCTTTTTATTTT-3' (CdRE-*E2F1*), respectively (the core sequence is underlined). The mutated sequences were 5'-AATTCGGCGGATTTTGCTGAATTTTGCG-3' (mCdRE-*HO-1*), 5'-AGTTTCATATTTGCTGAATATCAGTGTTTG-3' (mCdRE-*CCND1*), and 5'-zTTTGAACCTGATGCTGAATCTTTTTATTTT-3' (mCdRE-*E2F1*). EMSA was performed using in vitro-translated protein or the same amount of unprogrammed lysate (Promega) with LightShift Chemiluminescent EMSA Kits (Pierce). For competition experiments, a 100-fold molar excess of unlabeled CdRE was mixed with the biotin-labeled probe. The resulting protein-DNA complexes were analyzed by electrophoresis on a polyacrylamide gel, followed by chemiluminescent detection.

Anchorage-dependent and -independent growth assays. Anchorage-dependent cell proliferation was analyzed by a crystal violet assay as described previously (59). For anchorage-independent growth (57), 1×10^4 cells were plated on 6-cm plates containing a bottom layer of 0.6% low-melting-temperature agar in DMEM and a top layer of 0.3% agar in DMEM. Colonies were scored after 3 weeks of growth.

Animal experiments. Two days after implantation of estrogen pellets (E2, 0.36 mg/pellet, 60-day release) (Innovative Research of America), 1×10^7 tumor cells were injected into the abdominal mammary fat pad of 6-week-old female nude mice. When tumors reached the volume of approximately 100 mm³, we randomly allocated the mice to groups in which they received placebo or tamoxifen pellets (Innovative Research of America). Tumor growth was monitored by caliper measurements. Excised tumors were weighed, and portions were frozen in liquid nitrogen or fixed in 4% paraformaldehyde for further study.

Immunohistochemistry. Immunohistochemical staining was performed as described previously (59) using rabbit anti-ER α (Millipore), rabbit anti-ER β (Millipore), and rabbit anti-PES1 (Bethyl Laboratories) as primary antibodies.

Statistics. Differences among variables were assessed by χ^2 analysis or 2-tailed Student's *t* test. Estimation of disease-free and overall survival was performed using the Kaplan-Meier method, and differences between sur-



vival curves were determined with the log-rank test. Statistical calculations were performed using SPSS 13.0. *P* values of less than 0.05 were considered statistically significant.

Study approval. Animal studies were approved by the Institutional Animal Care Committee of Beijing Institute of Biotechnology. Breast cancer samples were obtained from Chinese PLA General Hospital, with the informed consent of patients and with institutional approval for experiments from Chinese PLA General Hospital and Beijing Institute of Biotechnology.

Acknowledgments

We thank Rong Li and Haian Fu for helpful discussions. This work was supported by Major State Basic Research Development

Program (2011CB504202 and 2012CB945100), National Natural Science Foundation (81072173, 30625035, and 30530320), and National Key Technologies R&D Program for New Drugs (2009ZX09301-002).

Received for publication December 30, 2011, and accepted in revised form May 31, 2012.

Address correspondence to: Qinong Ye, Department of Medical Molecular Biology, Beijing Institute of Biotechnology, 27 Tai-Ping Lu Rd., Beijing 100850, China. Phone: 8610.6818.0809; Fax: 8610.6824.8045; E-mail: yeqn66@yahoo.com.

1. Deroo BJ, Korach KS. Estrogen receptors and human disease. *J Clin Invest*. 2006;116(3):561–570.
2. Yager JD, Davidson NE. Estrogen carcinogenesis in breast cancer. *N Engl J Med*. 2006;354(3):270–282.
3. Thomas C, Gustafsson JA. The different roles of ER subtypes in cancer biology and therapy. *Nat Rev Cancer*. 2011;11(8):597–608.
4. Matthews J, Gustafsson JA. Estrogen signaling: a subtle balance between ER α and ER β . *Mol Interv*. 2003;3(5):281–292.
5. Helguero LA, Faulds MH, Gustafsson JA, Haldosen LA. Estrogen receptors α (ER α) and β (ER β) differentially regulate proliferation and apoptosis of the normal murine mammary epithelial cell line HC11. *Oncogene*. 2005;24(44):6605–6616.
6. Paruthiyil S, Parmar H, Kerekatte V, Cunha GR, Firestone GL, Leitman DC. Estrogen receptor β inhibits human breast cancer cell proliferation and tumor formation by causing a G2 cell cycle arrest. *Cancer Res*. 2004;64(1):423–428.
7. Omoto Y, Eguchi H, Yamamoto-Yamaguchi Y, Hayashi S. Estrogen receptor (ER) β 1 and ER β cx/ β 2 inhibit ER α function differently in breast cancer cell line MCF7. *Oncogene*. 2003;22(32):5011–5020.
8. Roger P, Sahla ME, Makela S, Gustafsson JA, Balder P, Rochefor H. Decreased expression of estrogen receptor β protein in proliferative preinvasive mammary tumors. *Cancer Res*. 2001;61(6):2537–2541.
9. Charpentier AH, et al. Effects of estrogen on global gene expression: identification of novel targets of estrogen action. *Cancer Res*. 2000;60(21):5977–5983.
10. Li J, et al. Down-regulation of pascadillo inhibits proliferation and tumorigenicity of breast cancer cells. *Cancer Sci*. 2009;100(12):2255–2260.
11. Zhang H, et al. The antibody preparation and expression of human Pascadillo. *Sci China C Life Sci*. 2007;50(3):298–304.
12. Nilsson S, Koehler KF, Gustafsson JA. Development of subtype-selective oestrogen receptor-based therapeutics. *Nat Rev Drug Discov*. 2011;10(10):778–792.
13. Shanley EK, Xu W. Selectively targeting estrogen receptors for cancer treatment. *Adv Drug Deliv Rev*. 2010;62(13):1265–1276.
14. Welboren WJ, Sweep FC, Span PN, Stunnenberg HG. Genomic actions of estrogen receptor α : what are the targets and how are they regulated? *Endocr Relat Cancer*. 2009;16(4):1073–1089.
15. Williams C, Edvardsson K, Lewandowski SA, Strom A, Gustafsson JA. A genome-wide study of the repressive effects of estrogen receptor β on estrogen receptor α signaling in breast cancer cells. *Oncogene*. 2008;27(7):1019–1032.
16. Bourdeau V, Deschenes J, Laperriere D, Aid M, White JH, Mader S. Mechanisms of primary and secondary estrogen target gene regulation in breast cancer cells. *Nucleic Acids Res*. 2008;36(1):76–93.
17. Chang EC, Frasor J, Komm B, Katzenellenbogen BS. Impact of estrogen receptor β on gene networks regulated by estrogen receptor α in breast cancer cells. *Endocrinology*. 2006;147(10):4831–4842.
18. Carroll JS, et al. Genome-wide analysis of estrogen receptor binding sites. *Nat Genet*. 2006;38(11):1289–1297.
19. Carroll JS, et al. Chromosome-wide mapping of estrogen receptor binding reveals long-range regulation requiring the forkhead protein FoxA1. *Cell*. 2005;122(1):33–43.
20. Lindberg MK, et al. Estrogen receptor (ER)- β reduces ER α -regulated gene transcription, supporting a “ying yang” relationship between ER α and ER β in mice. *Mol Endocrinol*. 2003;17(2):203–208.
21. Frasor J, Danes JM, Komm B, Chang KC, Lyttle CR, Katzenellenbogen BS. Profiling of estrogen up- and down-regulated gene expression in human breast cancer cells: insights into gene networks and pathways underlying estrogenic control of proliferation and cell phenotype. *Endocrinology*. 2003;144(10):4562–4574.
22. Coser KR, Chesnes J, Hur J, Ray S, Isselbacher KJ, Shioda T. Global analysis of ligand sensitivity of estrogen inducible and suppressible genes in MCF7/BUS breast cancer cells by DNA microarray. *Proc Natl Acad Sci U S A*. 2003;100(24):13994–13999.
23. Santen R, et al. Estrogen mediation of breast tumor formation involves estrogen receptor-dependent, as well as independent, genotoxic effects. *Ann N Y Acad Sci*. 2009;1155:132–140.
24. Foster JS, Henley DC, Ahamed S, Wimalasena J. Estrogens and cell-cycle regulation in breast cancer. *Trends Endocrinol Metab*. 2001;12(7):320–327.
25. Fox EM, Davis RJ, Shupnik MA. ER β in breast cancer—Onlooker, passive player, or active protector? *Steroids*. 2008;73(11):1039–1051.
26. Bretschneider N, Kangaspek S, Seifert M, Reid G, Gannon F, Denger S. E2-mediated cathepsin D (CTSD) activation involves looping of distal enhancer elements. *Mol Oncol*. 2008;2(2):182–190.
27. Eeckhoutte J, Carroll JS, Geistlinger TR, Torres-Arzayus MI, Brown M. A cell-type-specific transcriptional network required for estrogen regulation of cyclin D1 and cell cycle progression in breast cancer. *Genes Dev*. 2006;20(18):2513–2526.
28. Sikorski EM, Uo T, Morrison RS, Agarwal A. Pascadillo interacts with the cadmium response element of the human heme oxygenase-1 promoter in renal epithelial cells. *J Biol Chem*. 2006;281(34):24423–24430.
29. Shaaban AM, et al. Nuclear and cytoplasmic expression of ER β 1, ER β 2, and ER β 5 identifies distinct prognostic outcome for breast cancer patients. *Clin Cancer Res*. 2008;14(16):5228–5235.
30. Honma N, et al. Clinical importance of estrogen receptor- β evaluation in breast cancer patients treated with adjuvant tamoxifen therapy. *J Clin Oncol*. 2008;26(22):3727–3734.
31. Peng B, Lu B, Leygue E, Murphy LC. Putative functional characteristics of human estrogen receptor- β isoforms. *J Mol Endocrinol*. 2003;30(1):13–29.
32. Tateishi Y, et al. Ligand-dependent switching of ubiquitin-proteasome pathways for estrogen receptor. *EMBO J*. 2004;23(24):4813–4823.
33. Fan M, Park A, Nephew KP. CHIP (Carboxyl terminus of hsc70-interacting protein) promotes basal and geldanamycin-induced degradation of estrogen receptor- α . *Mol Endocrinol*. 2005;19(12):2901–2914.
34. Tateishi Y, et al. Turning off estrogen receptor β -mediated transcription requires estrogen-dependent receptor proteolysis. *Mol Cell Biol*. 2006;26(21):7966–7976.
35. Li L, Li Z, Howley PM, Sacks DB. E6AP and calmodulin reciprocally regulate estrogen receptor stability. *J Biol Chem*. 2006;281(4):1978–1985.
36. Picard N, et al. Phosphorylation of activation function-1 regulates proteasome-dependent nuclear mobility and E6-associated protein ubiquitin ligase recruitment to the estrogen receptor beta. *Mol Endocrinol*. 2008;22(2):317–330.
37. Chen GG, Vlantis AC, Zeng Q, van Hasselt CA. Regulation of cell growth by estrogen signaling and potential targets in thyroid cancer. *Curr Cancer Drug Targets*. 2008;8(5):367–377.
38. Bardin A, Boule N, Lazennec G, Vignon F, Pujol P. Loss of ER β expression as a common step in estrogen-dependent tumor progression. *Endocr Relat Cancer*. 2004;11(3):537–551.
39. Riggs BL, Hartmann LC. Selective estrogen-receptor modulators—mechanisms of action and application to clinical practice. *N Engl J Med*. 2003;348(7):618–629.
40. Jordan VC. Selective estrogen receptor modulation: concept and consequences in cancer. *Cancer Cell*. 2004;5(3):207–213.
41. Robertson JF. Fulvestrant (Faslodex)—how to make a good drug better. *Oncologist*. 2007;12(7):774–784.
42. Croxtall JD, McKeage K. Fulvestrant: a review of its use in the management of hormone receptor-positive metastatic breast cancer in postmenopausal women. *Drugs*. 2011;71(3):363–380.
43. Allende ML, Amsterdam A, Becker T, Kawakami K, Gaiano N, Hopkins N. Insertional mutagenesis in zebrafish identifies two novel genes, pascadillo and dead eye, essential for embryonic development. *Genes Dev*. 1996;10(24):3141–3155.
44. Lapiq YR, Fernandes CJ, Lau LF, Pestov DG. Physical and functional interaction between Pes1 and Bop1 in mammalian ribosome biogenesis. *Mol Cell*. 2004;15(1):17–29.
45. Killian A, et al. Inactivation of the RRB1-Pescadillo pathway involved in ribosome biogenesis induces chromosomal instability. *Oncogene*. 2004;23(53):8597–8602.
46. Grimm T, et al. Dominant-negative Pes1 mutants inhibit ribosomal RNA processing and cell proliferation via incorporation into the PeBoW-complex. *Nucleic Acids Res*. 2006;34(10):3030–3043.
47. Kinoshita Y, et al. Pascadillo, a novel cell cycle regulatory protein abnormally expressed in malignant cells. *J Biol Chem*. 2001;276(9):6656–6665.
48. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144(5):646–674.
49. Musgrove EA, Caldon CE, Barraclough J, Stone A, Sutherland RL. Cyclin D as a therapeutic target in cancer. *Nat Rev Cancer*. 2011;11(8):558–572.



50. Ray D, Terao Y, Christov K, Kaldis P, Kiyokawa H. Cdk2-null mice are resistant to ErbB-2-induced mammary tumorigenesis. *Neoplasia*. 2011; 13(5):439-444.
51. Meyer N, Penn LZ. Reflecting on 25 years with MYC. *Nat Rev Cancer*. 2008;8(12):976-990.
52. Ramsay RG, Gonda TJ. MYB function in normal and cancer cells. *Nat Rev Cancer*. 2008;8(7):523-534.
53. Stender JD, Frasor J, Komm B, Chang K, Kraus WL, Katzenellenbogen BS. Estrogen-regulated gene networks in human breast cancer cells: involvement of E2F1 in the regulation of cell proliferation. *Mol Endocrinol*. 2007;21(9):2112-2123.
54. Chen HZ, Tsai SY, Leone G. Emerging roles of E2Fs in cancer: an exit from cell cycle control. *Nat Rev Cancer*. 2009;9(11):785-797.
55. Maiorano D, Lutzmann M, Mechali M. MCM proteins and DNA replication. *Curr Opin Cell Biol*. 2006;18(2):130-136.
56. Bailis JM, Forsburg SL. MCM proteins: DNA damage, mutagenesis and repair. *Curr Opin Genet Dev*. 2004;14(1):17-21.
57. Ding L, et al. Human four-and-a-half LIM family members suppress tumor cell growth through a TGF- β -like signaling pathway. *J Clin Invest*. 2009;119(2):349-361.
58. He X, et al. c-Abl regulates estrogen receptor α transcription activity through its stabilization by phosphorylation. *Oncogene*. 2010;29(15):2238-2251.
59. Zhang H, et al. Stimulatory cross-talk between NFAT3 and estrogen receptor in breast cancer cells. *J Biol Chem*. 2005;280(52):43188-43197.
60. Han J, et al. Hepatitis B virus X protein and the estrogen receptor variant lacking exon 5 inhibit estrogen receptor signaling in hepatoma cells. *Nucleic Acids Res*. 2006;34(10):3095-3106.
61. Ding L, et al. Ligand-independent activation of estrogen receptor α by XBP-1. *Nucleic Acids Res*. 2003;31(18):5266-5274.